

CLAIMS

What is claimed is:

1. A method for treating or reducing the symptoms of gastroparesis in a patient comprising the step of:
administering metoclopramide or a pharmaceutically acceptable salt thereof
to a patient in need of gastroparesis treatment, wherein said metoclopramide
is in a pharmaceutically acceptable nasal formulation and administered
intranasally in a therapeutically effective amount so that one or more
symptom of gastroparesis is reduced.
2. The method of claim 1 wherein said daily dosage is between about 40 mg/day
to about 160 mg/day.
3. The method of claim 2 wherein said daily dosage is between about 40 mg/day
to about 80 mg/day.
4. The method of claim 2 wherein said daily dosage is about 40 mg/day.
5. The method of claim 2 wherein said daily dosage is about 80 mg/day.
6. The method of claim 1 wherein said daily dosage is divided into 3 or 4 equal
smaller doses and administered at equally spaced intervals within 24 hours.
7. The method of claim 1 wherein said daily dosage is divided into 3 or 4 equal
smaller doses and administered before meals and before bedtime.
8. The method of claim 6 or 7 wherein the smaller doses are about 10 mg each.
9. The method of claim 6 or 7 wherein the smaller doses are about 20 mg each.
10. The method of claim 1 wherein said daily dosage is between about 0.1 mg/kg
to about 2.5 mg/kg.
11. The method of claim 10, wherein said daily dosage is between about 0.6
mg/kg to about 1.2 mg/kg.

12. The method of claim 1 wherein the metoclopramide or pharmaceutically acceptable salt thereof is in an aqueous-based carrier.

13. The method of claim 1 wherein the metoclopramide or pharmaceutically acceptable salt thereof is in a sustained release formulation.

5

14. The method of claim 1 wherein the metoclopramide or pharmaceutically acceptable salt thereof is co-administered with one or more additional drugs.

15. The method of claim 1 wherein said dosage is administered for treating
10 gastroparesis caused by diabetes.

16. The method of claim 15 wherein said diabetes is selected from the group consisting of type 1 diabetes and type 2 diabetes.

17. The method of claim 1 wherein said dosage is administered for treating
15 gastroparesis caused by diabetes (including type 1 and type 2 diabetes), postviral syndromes, anorexia nervosa, surgery on the on the stomach or vagus nerve, medications, particularly anticholinergics and narcotics which slow contractions in the intestine, gastroesophageal reflux disease, smooth muscle disorders such as amyloidosis and scleroderma, nervous
20 system diseases (including abdominal migraine and Parkinson's disease), or metabolic disorders (including hypothyroidism).

18. The method of claim 1, 2, 3, 4 or 5 wherein said daily dosage is administered for about 1 weeks to about 8 weeks.

25

19. The method of claim 18 wherein said daily dosage is administered for about 5 weeks to about 8 weeks.

20. The method of claim 18 wherein said daily dosage is administered for about
30 6 weeks.

35